

(19)



Europäisches Patentamt

European Patent Office

Office européen des brevets



(11)

EP 0 939 082 A1

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
01.09.1999 Bulletin 1999/35

(51) Int. Cl.⁶: **C07K 14/375**, **A61K 38/16**,
A61K 35/84, **C12N 1/14**,
A01H 15/00

(21) Application number: **98403239.1**

(22) Date of filing: **21.12.1998**

(84) Designated Contracting States:
AT BE CH CY DE DK ES FI FR GB GR IE IT LI LU
MC NL PT SE
Designated Extension States:
AL LT LV MK RO SI

• **Okubo, Shigenobu**,
Gardenhills-Hisamatsu-102
Tsuchiura-shi, Ibaraki-ken (JP)
• **Hiroe, Eiji**
Himeji-shi, Hyogo-ken (JP)

(30) Priority: **16.01.1998 JP 1833498**

(71) Applicant:
Agaricus Laboratories Co., Ltd.
Chuo-ku, Tokyo (JP)

(74) Representative:
Clisci, Serge et al
S.A. FEDIT-LORIOT & AUTRES
CONSEILS EN PROPRIETE INDUSTRIELLE
38, Avenue Hoche
75008 Paris (FR)

(72) Inventors:
• **Hikichi, Manabu**
Sagamihara-shi, Kanagawa-ken (JP)

(54) **Protein polysaccharide 0041**

(57) The present invention relates to preparation of protein polysaccharide 0041 obtained by extracting a dried product of cultured mycelia of Agaricus blazei Murrill 0041 with hot water, and protein polysaccharide 0041 shows immunopotentiating antitumor activity when administered, so that it is useful as an antitumor agent against various tumors, and as an immunopotentiating substance against other diseases.

EP 0 939 082 A1

Description

DETAILED DESCRIPTION OF THE INVENTION

5 Technical Field to Which the Invention Belongs:

[0001] The present invention relates to protein polysaccharide 0041 with an immunopotentiating antitumor activity.

[0002] According to the present invention, protein polysaccharide 0041 has an immunopotentiating antitumor activity and thus high potency as an antitumor agent and an immunopotentiating substance for various tumors.

10

Prior Art:

[0003] Conventionally, fruit body of Agaricus blazei is regenerated and manufactured by artificially culturing its mycelia, and its dried fruit body is commercialized in a large amount as functional food.

15

[0004] The water extract of the dried fruit body, extracted with hot water, is drunk with the expectation of efficacy such as immunopotentialization.

Problem to be Solved by the Invention:

20

[0005] Preparation of the fruit body of Agaricus blazei requires a considerable day and time for regeneration of the fruit body from seed culture of the mycelia, and the productivity is low and its price for commercialization is consequently high.

[0006] There are a large number of Agaricus blazei mutants, and their efficacies of the biological activities are rather diverse and generally not uniform as a whole.

25

[0007] The object of the present invention is to search cultured mycelia of Agaricus blazei, not for the generation of the fruit body, but for a new strain of Agaricus blazei excellent in efficacy and stable productivity and to isolate a new immunopotentiating substance from the cultured mycelia.

Brief Description of the Drawings:

30

[0008]

Fig. 1 shows an ultraviolet absorption spectrum of protein polysaccharide 0041.

Fig. 2 shows an infrared absorption spectrum of protein polysaccharide 0041.

35

Fig. 3 shows an NMR spectrum of protein polysaccharide 0041.

Means to Solve the Problem:

40

[0009] The present inventors had isolated a large number of the strains with significantly-growing mycelia derived from many fruit bodies of Agaricus blazei cultivated at the place of origin in Brazil, then cultured each of them in a liquid medium, extracted the resulting mycelia with hot water, and examined the isolated substances for antitumor activity. As a result, the inventors recognized that only a certain strain has an immunopotentiating antitumor activity superior to that of other strains, thereby have performed the present invention.

45

[0010] In the present invention, the strain was designated Agaricus blazei Murrill 0041, and the substance with an immunopotentiating antitumor activity, prepared by the strain, was named protein polysaccharide 0041.

[0011] Agaricus blazei Murrill 0041 was deposited under FERM BP-6210 on December 18, 1997 with the National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology, Japan.

[0012] The microbial properties of Agaricus blazei Murrill 0041 are as follows:

50

1. It grows well in MY agar medium (30 g malt extract, 5 g yeast extract, 9 g agar, 1000 ml distilled water) at 27°C. The mycelia are white, grow densely, and are cream-colored when viewed through the cover of a Petri dish. The mycelia elongate radially, and some mycelia branch in two directions to elongate in a dendritic form, unlike the growth of di-nuclear hypha of a basidiomycete such as "shiitake" (Lentinula edodes) which elongate to form clump connections. In long-term culture, rhizomorph of the mycelia are partially formed, fruit bodies are not formed in the

55

Petri dish, and asexual spores are not observed.

2. It grows well in a potato/glucose agar medium as well and exhibits the same properties as in above 1.

3. Its mycelia grow well in shaking culture at 27°C in a liquid medium (50 g glucose, 5 g yeast extract, 30 g malt extract, adjusted to 1 liter with distilled water), and protein polysaccharide 0041 is formed in a large amount in the

mycelia.

[0013] Agaricus blazei Murrill 0041 is cultured for production of the mycelia.

[0014] Although culture of the mycelia may be carried out in either a solid or liquid medium, the cultivation in a liquid medium is more favorable for treatment of mycelia and is generally used.

[0015] The liquid medium may be any liquid media which are utilized for cultivation of the mycelia of the basidiomycota.

[0016] The medium for shaking culture is preferably a liquid medium prepared by mixing glucose (50 g/L), yeast extract (5.0 g/L), malt extract (30 g/L) and distilled water, and then adjusting the mixture to 1 liter volume with distilled water.

[0017] Culture is carried out aerobically in shaking culture, or under stirring and aeration, at about 27°C for 10 to 14 days, and the mycelia grow satisfactorily.

[0018] The cultured mycelia in the liquid medium are filtered under reduced pressure and then pressed in a filter press machine to give wet mycelia. The wet mycelia are dried in hot air to give dried mycelia.

[0019] The resulting dried mycelia are moved into a vessel and then mixed with distilled water which is 10 times the dried mycelia in volume, and refluxed for about 1 hour.

[0020] The obtained aqueous solution is subjected to press-filtration, and then further clarified by filtration using celite, followed by ultrafiltration treatment to remove low-molecular substances. The obtained retentate is concentrated to an about 1/2 to about 1/10 volume under reduced pressure, and the concentrate is mixed with a sufficient amount of ethanol and then allowed to stand overnight at room temperature to give a pale-brown precipitate. This precipitate is dried in hot air to give pale-brown powder of protein polysaccharide 0041.

[0021] The physicochemical properties of protein polysaccharide 0041 are as follows:

1. It is a protein polysaccharide.

2. The main saccharides of the polysaccharide are glucose and mannose.

3. It turns dark brown in phenol sulfate reaction which indicates 52.8 % of saccharide content.

4. It turns blue in Lowry-Folin reaction to indicate 27.2 % of protein content.

5. The amino acid constituents (%) obtained by measuring by an amino acid analyzer the hydrolysates of protein polysaccharide 0041 after hydrolysis with 6 N hydrochloric acid at 110°C for 12 hours in a sealed tube under reduced pressure are as follows: Asp 10.12, Thr 4.13, Ser 4.85, Glu 13.55, Gly 12.92, Ala 11.55, Val 7.44, Met 2.16, Ile 4.72, Leu 10.94, Tyr 2.68, Phe 3.91, Lys 4.62, His 2.05, Arg 2.40, Cys 0.60, and Pro 1.50.

6. Molecular weight: 85,000 to 105,000, determined by gel filtration chromatography.

7. Melting point: It turns brown at 245°C and dark brown at 275°C to 279°C; it decomposes at 275°C to 279°C.

8. Specific rotation: $[\alpha]_D^{21} = +318^\circ$ (0.33 % in water).

9. Solubility: Easily soluble in an aqueous 0.05 N NaOH solution and relatively soluble in water, and insoluble in methanol, ethanol, chloroform, benzene and hexane.

10. Color reaction: Positive in phenol/sulfate reaction, biuret reaction, Lowry-Folin reaction, and xanthoprotein reaction.

11. The pH value of the 1.58 % aqueous solution is 7.4.

12. The color of its powder is pale-brown.

13. UV absorption spectrum: As shown in Fig. 1.

14. IR absorption spectrum: As shown in Fig. 2.

15. NMR spectrum: As shown in Fig. 3.

[0022] Similar to protein polysaccharides obtained from other basidiomycetes, protein polysaccharide 0041 of the present invention is substantially innocuous and shows immunopotentiating antitumor activity in various administrations, so that it is useful as an antitumor agent and can be used as an effective ingredient for the production of an immunopotentiating composition.

[0023] The dosage of protein polysaccharide 0041 for oral administration is about 10 mg to about 1000 mg, preferably about 100 mg to about 200 mg/day/person.

Examples:

Example 1: Preparation of Protein Polysaccharide 0041

[0024] Agaricus blazei Murrill 0041 (FERM BP-6210) was cultured in a medium (prepared by dissolving 9 g agar, 30 g malt extract and 5 g yeast extract in distilled water, adjusting the total volume to 1 liter with distilled water, and sterilizing it in an autoclave at 121°C for 20 min.) in a Petri dish at 26°C in the dark for 20 days, and the white mycelia, which

grew well in a dendritic form, were stamped out with an about 12 mm cork borer to use as the seed culture.

[0025] Fifty grams of glucose, 5.0 g of yeast extract, and 30 g of malt extract were added to distilled water to prepare 1-liter liquid medium, and the liquid medium was poured into a 500-ml Sakaguchi flask in a volume of 100 ml/flask and then sterilized at 121°C for 45 minutes, and the above seed culture was inoculated into each flask and cultured at 27°C for 10 days. The culture from four flasks containing the cultured mycelia were combined and poured into a 20-liter culture tank (containing 12 liters of the same liquid medium as above) and cultured at 27°C under aeration at 0.5 vvm with stirring at 50 to 100 rpm for 14 days.

[0026] The cultural medium, 12 liters, was filtered under reduced pressure to give the mycelia. The mycelia were then pressed by a filter press machine to give the wet mycelia (water content of 65 to 75 %), and the wet mycelia were dried in hot air to give 6.0 g of the dried mycelia.

[0027] The dried mycelia were collected, and 50 g thereof was put into a 1000-ml round flask, and 500 ml of distilled water was added thereto. The resultant solution was heated under reflux for 1 hour, then cooled and filtered by press-filtration to give a filtrate, which was further filtered by using celite for clarification.

[0028] The clarified filtrate was subjected to ultrafiltration to remove substances of low molecular weights and the obtained retentate was concentrated to 100 ml under reduced pressure.

[0029] To 100 ml of the obtained concentrate, was added 400 ml of absolute ethanol. The mixture was allowed to stand overnight at room temperature to give precipitate, which was collected by filtration in vacuo and then dried in hot air whereby 4.8 g of protein polysaccharide 0041 was obtained.

Example 2: Immunopotentiating Antitumor Activity Test

[0030] The effect of protein polysaccharide 0041 against sarcoma 180 tumor was examined.

[0031] Mice (30 g male ICR) were inoculated with the tumor (0.1 ml subcutaneous inoculation onto the back of a mouse: 3×10^6 tumor cells). Twenty four hours later, an aqueous solution prepared by dissolving pale-brown protein polysaccharide 0041 obtained in Example 1 in physiological saline (1 mg/ml) was administered intraperitoneally into the mice as one group 10 times for 10 days at a dose of 10 mg/kg/day. Otherwise, physiological saline only was administered into a control group in the test. Twenty days after, i.e., on the day of final dosage, the tumor nodes were removed from the group given the sample and the control group, and the weight of each tumor node was measured. The rate of inhibition is calculated using the following equation:

$$\text{Rate of inhibition (\%)} = (C - T)/C \times 100$$

where T is the average weight of tumor nodes from the group given the sample and C is that from the control group.

[0032] The results are as follows:

Group	Tumor average weight (g)	Degree of tumor inhibition (%)	Complete disappearance
Control group	12.64		0/10 (Two of ten mice died)
Group given protein polysaccharide 0041	3.60	71.5	4/10

Claims

1. Protein polysaccharide 0041 having the following physicochemical properties:

- (1) it is a protein polysaccharide;
- (2) the main saccharides of the polysaccharide are glucose and mannose.
- (3) molecular weight: 85,000 to 105,000, determined by gel filtration chromatography;
- (4) melting point: it turns brown at 245°C and dark brown at 275 to 279°C; it decomposes at 275 to 279°C;
- (5) specific rotation: $[\alpha]_D^{21} = +318^\circ$ (0.33 % in water);
- (6) solubility: it dissolves easily in an aqueous 0.05 N NaOH solution, and dissolves relatively in water and is insoluble in methanol, ethanol, chloroform, benzene and hexane;
- (7) color reaction: positive in phenol/sulfate reaction, biuret reaction, Lowry-Folin reaction, and xanthoprotein

reaction;

(8) the pH value of the 1.58 % aqueous solution is 7.4; and

(9) the color of its powder is pale-brown.

- 5 2. Protein polysaccharide 0041 obtained by extracting a dried product from cultured mycelia of Agaricus blazei Murrill 0041 with hot water.
3. An antitumor agent comprising protein polysaccharide 0041 as the potential ingredient.
- 10 4. An immunopotentiating composition comprising protein polysaccharide 0041.
5. A process for producing protein polysaccharide 0041, which comprises culturing Agaricus blazei Murrill 0041, drying the obtained mycelia from necessity, subjecting it to extraction treatment, and isolating protein polysaccharide 0041.
- 15 6. Agaricus blazei Murrill 0041, FERM BP-6210.

20

25

30

35

40

45

50

55

FIG. 1





FIG. 2






FIG. 3



European Patent
Office

EUROPEAN SEARCH REPORT

Application Number
EP 98 40 3239

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.6)
X	PATENT ABSTRACTS OF JAPAN vol. 014, no. 509 (C-0776), 7 November 1990 & JP 02 211847 A (IWADE KINGAKU KENKYUSHO;KK;OTHERS: 01), 23 August 1990 * abstract *	1-5	C07K14/375 A61K38/16 A61K35/84 C12N1/14 A01H15/00
X	PATENT ABSTRACTS OF JAPAN vol. 014, no. 270 (C-0727), 12 June 1990 & JP 02 078630 A (NICHIREI CORP;OTHERS: 04), 19 March 1990 * abstract *	1-5	
			TECHNICAL FIELDS SEARCHED (Int.Cl.6)
			A61K C12N
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 22 April 1999	Examiner Rempp, G
CATEGORY OF CITED DOCUMENTS X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			

EPO FORM 1503/03B2 (P4/C01)